This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images, please do not report the images to the Image Problems Mailbox.

COMPARISON OF WAVELENGTH DISTRIBUTION OF UV SOURCES



FIG. 1

FIG 2A SE500660



FIG. 2B SE500660

SOLAR SIMULATED LIGHT ACTIVATES NF-κB IN HUMAN SKIN *IN VIVO*



TOBACO 5590660

FIG. 3A

SOLAR-SIMULATED UV INDUCTION OF COLLAGENASE IN HUMAN SKIN *IN VIVO*



TOOLO'SESOOGGO

FIG. 3B

DOSE DEPENDENCE FOR SOLAR-STIMULATED UV INDUCTION OF 92kDa GELATINASE ACTIVITY IN HUMAN SKIN *IN VIVO*



FIG. 3C

SOLAR SIMULATED UV REDUCES TYPE I PROCOLLAGEN IN HUMAN SKIN *IN VIVO*



T09020 "SES00660

UVB & LONG UVA INDUCE COLLAGENASE (MMP-1) mRNA IN HUMAN SKIN *IN VIVO*

FIG. 4A



FIG. 4B

CHARACTERISTICS OF UV FILTERS USED TO DETERMINE WAVELENGTH DEPENDENCE OF COLLAGENASE INDUCTION IN HUMAN SKIN *IN VIVO*



SPECTRAL OUPUT OF SOLAR SIMULATOR WITH MONOCHROMATOR: DETERMINING WHAT WAVELENGTHS CAUSE PHOTOAGING



FIG. 5

FIG. 6A


92kDa GELATINASE ACTIVITY INDUCED BY UVB (300-320nm) & LONG WAVELENGTH UVA1 (380nm): MONOCHROMATOR
92kDa GELATINASE ACTIVITY (FOLD INDUCTION)
20
15
10
5
0
.5J/cm2
5J/cm2
5J/cm2
5J/cm2
5J/cm2
300nm n=6
320nm n=5
340nm n=6
360nm n=6
380nm n=6


UV ACTION SPECTRUM FOR INDUCTION OF
92kDa GELATINASE (MMP-9) ACTIVITY
92kDa Gelatinase Activity
(Fold Induction)
12
11
10
9
8
7
6
5
4
3
2
1
285-305nm
305-325nm
365-385nm
345-365nm
325-345nm
385-425nm
0
2
4
6
8
10
UV Exposure (J/cm2)

FIG. 6B

FIG. 6C

RELATIVE EFFECTIVENESS OF UV WAVELENGTH TO INDUCE 92 kDa GELATINASE ACTIVITY IN HUMAN SKIN *IN VIVO*



FIG. 7

EFFECTIVE CONTRIBUTION OF UV WAVELENGTHS TO INDUCTION OF 92kDa GELATINASE ACTIVITY BY SOLAR-SIMULATED UV IN HUMAN SKIN *IN VIVO*



1MED 2MED 3MED

FIG. 8


SUNLIGHT VARIABILITY
UVB
UVA2
UVA1
Noon
Early morning /
Late afternooon
Irradiance (watts/nm/cm²)
10-4
10-5
10-6
300
320
340
360
380
400
Wavelength (nm)

FIG. 9

ABSORPTION SPECTRA OF PARSOL 1789 & MCX IN ABSOLUTE ETHANOL



FIG. 10

UVA1 WAVELENGTH DEPENDENCE FOR INDUCTION OF COLLAGENASE mRNA IN HUMAN SKIN *IN VIVO*



UVB/A2 Wavelength Dependence for Induction of Collagenase mRNA in Human skin in vivo

NO UV	300	310	320	330	340	350	360	
3.53E+02	1.05E+05	6.70E+04	2.88E+02	2.84E+02	5.05E+02	3.80E+02	3.09E+02	mean
1.31E+02	2.72E+04	8.96E+04	6.34E+01	5.71E+01	2.86E+02	1.13E+02	1.38E+02	sem

FIG. 11